

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

**Figure 1A: Retention of Activity by PEGylated Candida Uricase**



**Figure 1B: Retention of Activity by PEGylated Candida Uricase**




Figure 2A: Retention of Activity by PEGylated Porcine Uricase




Figure 2B: Retention of Activity by PEGylated Porcine Uricase




Figure 3A: Retention of Activity by PEGylated PBC Uracase





Figure 3B: Retention of Activity by PEGylated PBC Uricase



**Figure 4A:**  
**Retention of Activity by PEGylated Uricozyme® (*A. flavus* Uricase)**






Figure 4B:  
Retention of Activity by PEGylated Uricozyme® (*A. flavus* Uricase)




Figure 5A: Retention of Activity by PEGylated Soybean Uricase




Figure 5B: Retention of Activity by PEGylated Soybean Uricase






Figure 6: Deduced amino acid sequences of Pig-Baboon Chimeric (PBC) uricase, PBC uricase that is truncated at the amino and carboxyl terminals (PBC-NT-CT) and Porcine uricase containing the mutations R291K and T301S (PKS Uracase) (SEQ ID NO:3), compared with the porcine sequence (SEQ ID NO: 1) and baboon sequence (SEQ ID NO: 2)

Porcine	MAHYRNDYKK NDEVEFVRTG YGKDMIKVLH IQRDGKYHSI	40
PBC	porcine sequence 1-225 →	40
PBC-NT-CT	porcine sequence 1-219 →	34
PKS	porcine sequence 1-288 →	40
Baboon	MADYHNNYKK NDELEFVRTG YGKDMVKVLH IQRDGKYHSI	40
Porcine	KEVATSVQLT LSSKKDYLHG DNSDVIPTDT IKNTVNVLAK	80
PBC	porcine sequence →	80
PBC-NT-CT	porcine sequence →	74
PKS	porcine sequence →	80
Baboon	KEVATSVQLT LSSKKDYLHG DNSDIIPTDT IKNTVHVLAKE	80
Porcine	FKGIKSIETF AVTICEHFLS SFKHVIRAQV YVEEVPWKRF	120
PBC	porcine sequence →	120
PBC-NT-CT	porcine sequence →	114
PKS	porcine sequence →	120
Baboon	FKGIKSIEAF GVNICEYFLS SFNHVIRAQV YVEEIPWKRL	120
Porcine	EKNGVKHVHA FIYTPTGTHF CEVEQIRNGP PVIHSGIKDL	160
PBC	porcine sequence →	160
PBC-NT-CT	porcine sequence →	154
PKS	porcine sequence →	160
Baboon	EKNGVKHVHA FIHTPTGTHF CEVEQLRSGP PVIHSGIKDL	160
Porcine	KVLKTTQSGF EGFIKDQFTT LPEVKDRCFA TQVYCKWRYH	200
PBC	porcine sequence →	200
PBC-NT-CT	porcine sequence →	194
PKS	porcine sequence →	200
Baboon	KVLKTTQSGF EGFIKDQFTT LPEVKDRCFA TQVYCKWRYH	200
Porcine	QGRDVFDEAT WDTVRSIVLQ KFAGPYDKGE YSPSVQKTLY	240
PBC	porcine sequence → ← baboon sequence	240
PBC-NT-CT	porcine sequence → ← baboon sequence	234
PKS	porcine sequence →	240
Baboon	QCRDVFDEAT WGTIRDLVLE KFAGPYDKGE YSPSVQKTLY	240
Porcine	DIQVLTLGQV PEIEDMEISL PNIHYLNIDM SKMGLINKEE	280
PBC	baboon sequence →	280
PBC-NT-CT	baboon sequence →	274
PKS	porcine sequence →	280
Baboon	DIQVLSLSRV PEIEDMEISL PNIHYFNIDM SKMGLINKEE	280
Porcine	VLLPLDNPYG RITGTVKRKL TSRL	304
PBC	baboon sequence →	304
PBC-NT-CT	baboon sequence →	295
PKS	porcine   ← baboon →	304
Baboon	VLLPLDNPYG KITGTVKRKL SSRL	304


**Figure 7: Serum Uricase Activity 24 Hours after Each PEG-Uricase Injection, Relative to the First Injection**



**Figure 8: Inverse Relationship between Serum PEG-Uricase Activity and Uric Acid Levels in the Serum and Urine of a Uricase-Deficient Mouse**



**Figure 9: Decreased Severity of Urine-Concentrating Defect  
in Uricase-Deficient Mice Treated with PEG-Uricase**



**Figure 10: Decreased Severity of Nephrogenic Diabetes Insipidus  
in Uricase-Deficient Mice Treated with PEG-Uricase**







Figure 11:

Decreased Severity of Uric Acid-Induced Nephropathy after Treatment with PEG-Uricase, as Visualized by Magnetic Resonance Microscopy



**Figure 12: Clearance from the Circulation of BALB/c Mice of PBC Uricase Tetramer and Octamer Coupled to 5-6 Strands of 10-kDa PEG/Subunit**

